

A Logical Approach to Discrete Math

Mathematical Induction

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$P.1 :$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$P.2 :$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2 \quad \text{true}$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2 \quad \text{true}$$

$P.3 :$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2 \quad \text{true}$$

$$P.3 : 1 + 3 + 5 = 3^2$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2 \quad \text{true}$$

$$P.3 : 1 + 3 + 5 = 3^2 \quad \text{true}$$

A Logical Approach to Discrete Math

Mathematical Induction

$$P.n : (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$P.1 : 1 = 1^2 \quad \text{true}$$

$$P.2 : 1 + 3 = 2^2 \quad \text{true}$$

$$P.3 : 1 + 3 + 5 = 3^2 \quad \text{true}$$

How can you prove $P.n$ is true for all $n \geq 1$, not just $1 \leq n \leq 3$?

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

...

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

...

Conclusion: $P.n$ is true for all $n \geq 1$.

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

...

Conclusion: $P.n$ is true for all $n \geq 1$.

Proving $P.1$ is called the base case.

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

...

Conclusion: $P.n$ is true for all $n \geq 1$.

Proving $P.1$ is called the base case.

Proving $P.n \Rightarrow P(n + 1)$ by deduction is called the induction case.

A Logical Approach to Discrete Math

Idea: Suppose you could prove $P.n \Rightarrow P(n + 1)$ in general.

Then you could use (3.77) Modus ponens $p \wedge (p \Rightarrow q) \Rightarrow q$ as follows:

First prove $P.1$.

Then,

$$P.1 \wedge (P.1 \Rightarrow P.2) \Rightarrow P.2$$

$$P.2 \wedge (P.2 \Rightarrow P.3) \Rightarrow P.3$$

$$P.3 \wedge (P.3 \Rightarrow P.4) \Rightarrow P.4$$

...

Conclusion: $P.n$ is true for all $n \geq 1$.

Proving $P.1$ is called the base case.

Proving $P.n \Rightarrow P(n + 1)$ by deduction is called the induction case.

The antecedent $P.n$, which you assume, is called the inductive hypothesis.

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2 \\ = & \langle \text{Base case, } n = 1 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2 \\ = & \langle \text{Base case, } n = 1 \rangle \\ & (\sum i \mid 1 \leq i \leq 1 : 2 \cdot i - 1) = 1^2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$= \langle \text{Base case, } n = 1 \rangle$$

$$(\sum i \mid 1 \leq i \leq 1 : 2 \cdot i - 1) = 1^2$$

$$= \langle \text{Math} \rangle$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$= \langle \text{Base case, } n = 1 \rangle$$

$$(\sum i \mid 1 \leq i \leq 1 : 2 \cdot i - 1) = 1^2$$

$$= \langle \text{Math} \rangle$$

$$2 \cdot 1 - 1 = 1$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$$

$$= \langle \text{Base case, } n = 1 \rangle$$

$$(\sum i \mid 1 \leq i \leq 1 : 2 \cdot i - 1) = 1^2$$

$$= \langle \text{Math} \rangle$$

$$2 \cdot 1 - 1 = 1$$

$$= \langle \text{Math} \rangle$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2 \\ = & \langle \text{Base case, } n = 1 \rangle \\ & (\sum i \mid 1 \leq i \leq 1 : 2 \cdot i - 1) = 1^2 \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 1 - 1 = 1 \\ = & \langle \text{Math} \rangle \\ & 1 = 1 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1)$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n^2 + 2(n + 1) - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n^2 + 2(n + 1) - 1 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n^2 + 2(n + 1) - 1 \\ = & \langle \text{Math} \rangle \\ & n^2 + 2n + 1 \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n^2 + 2(n + 1) - 1 \\ = & \langle \text{Math} \rangle \\ & n^2 + 2n + 1 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $(\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) = (n + 1)^2$

assuming $(\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) = n^2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 1 \leq i \leq n + 1 : 2 \cdot i - 1) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 1 \leq i \leq n : 2 \cdot i - 1) + 2(n + 1) - 1 \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n^2 + 2(n + 1) - 1 \\ = & \langle \text{Math} \rangle \\ & n^2 + 2n + 1 \\ = & \langle \text{Math} \rangle \\ & (n + 1)^2 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 2^i) = 2^0 - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 2^i) = 2^0 - 1 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 2^i) = 2^0 - 1 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 2^i) = 2^0 - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 2^i) = 2^0 - 1 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 2^i) = 2^0 - 1 \\ = & \langle (8.13) \text{ Empty range rule, and math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 2^i) = 2^0 - 1 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 2^i) = 2^0 - 1 \\ = & \langle (8.13) \text{ Empty range rule, and math} \rangle \\ & 0 = 0 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$(\sum i \mid 0 \leq i < n + 1 : 2^i)$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & 2^n - 1 + 2^n \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & 2^n - 1 + 2^n \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & 2^n - 1 + 2^n \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & 2^n - 1 + 2^n \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n - 1 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 2^i) = 2^{n+1} - 1$

assuming $(\sum i \mid 0 \leq i < n : 2^i) = 2^n - 1$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 2^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 2^i) + 2^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & 2^n - 1 + 2^n \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n - 1 \\ = & \langle \text{Math} \rangle \\ & 2^{n+1} - 1 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 3^i) = (3^0 - 1)/2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 3^i) = (3^0 - 1)/2 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 3^i) = (3^0 - 1)/2 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 3^i) = (3^0 - 1)/2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 3^i) = (3^0 - 1)/2 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 3^i) = (3^0 - 1)/2 \\ = & \langle (8.13) \text{ Empty range rule, and math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} & (\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2 \\ = & \langle \text{Base case, } n = 0 \rangle \\ & (\sum i \mid 0 \leq i < 0 : 3^i) = (3^0 - 1)/2 \\ = & \langle \text{Math} \rangle \\ & (\sum i \mid \text{false} : 3^i) = (3^0 - 1)/2 \\ = & \langle (8.13) \text{ Empty range rule, and math} \rangle \\ & 0 = 0 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$(\sum i \mid 0 \leq i < n + 1 : 3^i)$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \\ & (3^n - 1 + 2 \cdot 3^n)/2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \\ & (3^n - 1 + 2 \cdot 3^n)/2 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \\ & (3^n - 1 + 2 \cdot 3^n)/2 \\ = & \langle \text{Math} \rangle \\ & (3 \cdot 3^n - 1)/2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \\ & (3^n - 1 + 2 \cdot 3^n)/2 \\ = & \langle \text{Math} \rangle \\ & (3 \cdot 3^n - 1)/2 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$ for $n \geq 0$

Induction case

Prove $(\sum i \mid 0 \leq i < n + 1 : 3^i) = (3^{n+1} - 1)/2$

assuming $(\sum i \mid 0 \leq i < n : 3^i) = (3^n - 1)/2$

as the inductive hypothesis.

$$\begin{aligned} & (\sum i \mid 0 \leq i < n + 1 : 3^i) \\ = & \langle \text{Split off last term} \rangle \\ & (\sum i \mid 0 \leq i < n : 3^i) + 3^n \\ = & \langle \text{Inductive hypothesis} \rangle \\ & (3^n - 1)/2 + 3^n \\ = & \langle \text{Math, common denominator} \rangle \\ & (3^n - 1 + 2 \cdot 3^n)/2 \\ = & \langle \text{Math} \rangle \\ & (3 \cdot 3^n - 1)/2 \\ = & \langle \text{Math} \rangle \\ & (3^{n+1} - 1)/2 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

$$2n + 1 < 2^n$$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

$$\begin{aligned} & 2n + 1 < 2^n \\ = & \langle \text{Base case, } n = 3 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

$$2n + 1 < 2^n$$

$$= \langle \text{Base case, } n = 3 \rangle$$

$$2 \cdot 3 + 1 < 2^3$$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

$$2n + 1 < 2^n$$

$$= \langle \text{Base case, } n = 3 \rangle$$

$$2 \cdot 3 + 1 < 2^3$$

$$= \langle \text{Math} \rangle$$

A Logical Approach to Discrete Math

Prove $2n + 1 < 2^n$ for $n \geq 3$

Proof

Base case

$$2n + 1 < 2^n$$

$$= \langle \text{Base case, } n = 3 \rangle$$

$$2 \cdot 3 + 1 < 2^3$$

$$= \langle \text{Math} \rangle$$

$$7 < 8 \quad //$$

A Logical Approach to Discrete Math

Induction case

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$2^{n+1}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$2^{n+1} \\ = \langle \text{Math} \rangle$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

2^{n+1}

= $\langle \text{Math} \rangle$

$2 \cdot 2^n$

> $\langle \text{Inductive hypothesis} \rangle$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \\ > & \langle \text{Inductive hypothesis} \rangle \\ & 2 \cdot (2n + 1) \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \\ > & \langle \text{Inductive hypothesis} \rangle \\ & 2 \cdot (2n + 1) \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \\ > & \langle \text{Inductive hypothesis} \rangle \\ & 2 \cdot (2n + 1) \\ = & \langle \text{Math} \rangle \\ & 2(n + 1) + 1 + 2n - 1 \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n + 1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \\ > & \langle \text{Inductive hypothesis} \rangle \\ & 2 \cdot (2n + 1) \\ = & \langle \text{Math} \rangle \\ & 2(n + 1) + 1 + 2n - 1 \\ > & \langle 2n - 1 \text{ is positive for } n \geq 3 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Induction case

Prove $2(n+1) + 1 < 2^{n+1}$

assuming $2n + 1 < 2^n$

as the inductive hypothesis.

$$\begin{aligned} & 2^{n+1} \\ = & \langle \text{Math} \rangle \\ & 2 \cdot 2^n \\ > & \langle \text{Inductive hypothesis} \rangle \\ & 2 \cdot (2n + 1) \\ = & \langle \text{Math} \rangle \\ & 2(n+1) + 1 + 2n - 1 \\ > & \langle 2n - 1 \text{ is positive for } n \geq 3 \rangle \\ & 2(n+1) + 1 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Example of a proof by induction. Consider a currency consisting of 2-cent and 5-cent coins. Show that any amount above 3 cents can be represented using these coins.

We write $P.n$ in English as

$P.n$: Some bag of 2-cent and 5-cent coins has the sum n .

Our task is to prove $(\forall n \mid 4 \leq n : P.n)$.

A Logical Approach to Discrete Math

Prove any amount more than 3 cents is possible using only 2-cent and 5-cent coins.

A Logical Approach to Discrete Math

Prove any amount more than 3 cents is possible using only 2-cent and 5-cent coins.

Base case

A Logical Approach to Discrete Math

Prove any amount more than 3 cents is possible using only 2-cent and 5-cent coins.

Base case

The base case is $n=4$.

A Logical Approach to Discrete Math

Prove any amount more than 3 cents is possible using only 2-cent and 5-cent coins.

Base case

The base case is $n=4$.

Must prove that you can make 4 cents using only 2-cent and 5-cent coins.

A Logical Approach to Discrete Math

Prove any amount more than 3 cents is possible using only 2-cent and 5-cent coins.

Base case

The base case is $n=4$.

Must prove that you can make 4 cents using only 2-cent and 5-cent coins.

Use two 2-cent coins. //

A Logical Approach to Discrete Math

Induction case

A Logical Approach to Discrete Math

Induction case

Must prove

A Logical Approach to Discrete Math

Induction case

Must prove

" $n+1$ cents is possible with 2-cent and 5-cent coins"

A Logical Approach to Discrete Math

Induction case

Must prove

" $n+1$ cents is possible with 2-cent and 5-cent coins"

assuming

A Logical Approach to Discrete Math

Induction case

Must prove

" $n+1$ cents is possible with 2-cent and 5-cent coins"

assuming

" n cents is possible with 2-cent and 5-cent coins"

A Logical Approach to Discrete Math

Induction case

Must prove

" $n+1$ cents is possible with 2-cent and 5-cent coins"

assuming

" n cents is possible with 2-cent and 5-cent coins"

as the inductive hypothesis.

A Logical Approach to Discrete Math

Case I

A Logical Approach to Discrete Math

Case I

You have n cents with at least one 5-cent coin. Remove

A Logical Approach to Discrete Math

Case I

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins.

A Logical Approach to Discrete Math

Case I

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins,

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins, they must all be 2-cent coins. Because the amount must

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins, they must all be 2-cent coins. Because the amount must be more than three cents, you must have at least two 2-

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins, they must all be 2-cent coins. Because the amount must be more than three cents, you must have at least two 2-cent coins. Remove two 2-cent coins and replace them

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins, they must all be 2-cent coins. Because the amount must be more than three cents, you must have at least two 2-cent coins. Remove two 2-cent coins and replace them with one 5-cent coin. Now you have $n+1$ cents with only

A Logical Approach to Discrete Math

Case 1

You have n cents with at least one 5-cent coin. Remove one 5-cent coin and replace it with three 2-cent coins. Now you have $n+1$ cents with only 2-cent and 5-cent coins.

Case 2

You have no five cent coins. If you have no 5-cent coins, they must all be 2-cent coins. Because the amount must be more than three cents, you must have at least two 2-cent coins. Remove two 2-cent coins and replace them with one 5-cent coin. Now you have $n+1$ cents with only 2-cent and 5-cent coins. //

A Logical Approach to Discrete Math

(12.11) **Definition, b to the power n :**

$$b^0 = 1$$

$$b^{n+1} = b \cdot b^n \quad \text{for } n \geq 0$$

(12.12) **b to the power n :**

$$b^0 = 1$$

$$b^n = b \cdot b^{n-1} \quad \text{for } n \geq 1$$

(12.13) **Definition, factorial:**

$$0! = 1$$

$$n! = n \cdot (n - 1)! \quad \text{for } n > 0$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$n! = (\prod i \mid 1 \leq i \leq n : i)$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \\ 0! &= (\prod i \mid 1 \leq i \leq 0 : i) \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \\ 0! &= (\prod i \mid 1 \leq i \leq 0 : i) \\ &= \langle (12.13) \text{ and math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \\ 0! &= (\prod i \mid 1 \leq i \leq 0 : i) \\ &= \langle (12.13) \text{ and math} \rangle \\ 1 &= (\prod i \mid \text{false} : i) \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \\ 0! &= (\prod i \mid 1 \leq i \leq 0 : i) \\ &= \langle (12.13) \text{ and math} \rangle \\ &= 1 = (\prod i \mid \text{false} : i) \\ &= \langle (8.13) \text{ Empty range rule} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Proof

Base case

$$\begin{aligned} n! &= (\prod i \mid 1 \leq i \leq n : i) \\ &= \langle \text{Base case, } n = 0 \rangle \\ 0! &= (\prod i \mid 1 \leq i \leq 0 : i) \\ &= \langle (12.13) \text{ and math} \rangle \\ &= 1 = (\prod i \mid \text{false} : i) \\ &= \langle (8.13) \text{ Empty range rule} \rangle \\ &= 1 = 1 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n + 1)! = (\prod i \mid 1 \leq i \leq n + 1 : i)$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n + 1)! = (\prod i \mid 1 \leq i \leq n + 1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$(\prod i \mid 1 \leq i \leq n+1 : i)$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \\ & (\prod i \mid 1 \leq i \leq n : i) \cdot (n+1) \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \\ & (\prod i \mid 1 \leq i \leq n : i) \cdot (n+1) \\ = & \langle \text{Inductive hypothesis} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \\ & (\prod i \mid 1 \leq i \leq n : i) \cdot (n+1) \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n! \cdot (n+1) \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \\ & (\prod i \mid 1 \leq i \leq n : i) \cdot (n+1) \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n! \cdot (n+1) \\ = & \langle (12.13 \text{ with } n := n+1, \text{ which is } (n+1)! = (n+1) \cdot n! \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove $n! = (\prod i \mid 1 \leq i \leq n : i)$ for $n \geq 0$

Induction case

Prove $(n+1)! = (\prod i \mid 1 \leq i \leq n+1 : i)$

assuming $n! = (\prod i \mid 1 \leq i \leq n : i)$

as the inductive hypothesis.

$$\begin{aligned} & (\prod i \mid 1 \leq i \leq n+1 : i) \\ = & \langle \text{Split off last term} \rangle \\ & (\prod i \mid 1 \leq i \leq n : i) \cdot (n+1) \\ = & \langle \text{Inductive hypothesis} \rangle \\ & n! \cdot (n+1) \\ = & \langle (12.13 \text{ with } n := n+1, \text{ which is } (n+1)! = (n+1) \cdot n! \rangle \\ & (n+1)! \quad // \end{aligned}$$

A Logical Approach to Discrete Math

The Golden Ratio


A Logical Approach to Discrete Math

The Golden Ratio




A Logical Approach to Discrete Math

The Golden Ratio




A Logical Approach to Discrete Math

The Golden Ratio



A Logical Approach to Discrete Math


The Golden Ratio



The golden ratio is $\phi = A/B$

A Logical Approach to Discrete Math


The Golden Ratio



The golden ratio is $\phi = A/B$
By definition

A Logical Approach to Discrete Math

The Golden Ratio




The golden ratio is $\phi = A/B$

By definition

$$\frac{A}{B} = \frac{A+B}{A}$$

A Logical Approach to Discrete Math

The Golden Ratio



The golden ratio is $\phi = A/B$


By definition

$$\frac{A}{B} = \frac{A+B}{A}$$

$$\frac{A}{B} = 1 + \frac{B}{A}$$

A Logical Approach to Discrete Math

The Golden Ratio



The golden ratio is $\phi = A/B$

By definition


$$\frac{A}{B} = \frac{A+B}{A}$$

$$\frac{A}{B} = 1 + \frac{B}{A}$$

$$\frac{A}{B} = 1 + \frac{1}{A/B}$$

A Logical Approach to Discrete Math

The Golden Ratio



The golden ratio is $\phi = A/B$

By definition

$$\frac{A}{B} = \frac{A+B}{A}$$

$$\frac{A}{B} = 1 + \frac{B}{A}$$

$$\frac{A}{B} = 1 + \frac{1}{A/B}$$

$$\phi = 1 + \frac{1}{\phi}$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

$$\phi^2 = \phi + 1$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

$$\phi^2 = \phi + 1$$

$$\phi^2 - \phi - 1 = 0$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

$$\phi^2 = \phi + 1$$

$$\phi^2 - \phi - 1 = 0$$

$$\phi = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-1)}}{2(1)}$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

$$\phi^2 = \phi + 1$$

$$\phi^2 - \phi - 1 = 0$$

$$\phi = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-1)}}{2(1)}$$

$$\phi = \frac{1 \pm \sqrt{5}}{2}$$

A Logical Approach to Discrete Math

$$\phi = 1 + \frac{1}{\phi}$$

$$\phi^2 = \phi + 1$$

$$\phi^2 - \phi - 1 = 0$$

$$\phi = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-1)}}{2(1)}$$

$$\phi = \frac{1 \pm \sqrt{5}}{2}$$

$$\phi = \frac{1 + \sqrt{5}}{2} \quad \hat{\phi} = \frac{1 - \sqrt{5}}{2}$$

A Logical Approach to Discrete Math

The Fibonacci sequence

0	1	1	2	3	5	8
F_0	F_1	F_2	F_3	F_4	F_5	F_6

A Logical Approach to Discrete Math

(12.14) **Definition, Fibonacci:**

$$F_0 = 0, \quad F_1 = 1$$

$$F_n = F_{n-1} + F_{n-2} \quad \text{for } n > 1$$

(12.14.1) **Definition, Golden Ratio:** $\phi = (1 + \sqrt{5})/2 \approx 1.618$ $\hat{\phi} = (1 - \sqrt{5})/2 \approx -0.618$

$$(12.15) \quad \phi^2 = \phi + 1 \quad \text{and} \quad \hat{\phi}^2 = \hat{\phi} + 1$$

$$(12.16) \quad F_n \leq \phi^{n-1} \quad \text{for } n \geq 1$$

$$(12.16.1) \quad \phi^{n-2} \leq F_n \quad \text{for } n \geq 1$$

$$(12.17) \quad F_{n+m} = F_m \cdot F_{n+1} + F_{m-1} \cdot F_n \quad \text{for } n \geq 0 \text{ and } m \geq 1$$

A Logical Approach to Discrete Math

To prove Fibonacci theorems
there are two base cases
and two inductive hypotheses.

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$
$$= \langle \text{Base case, } n = 1 \rangle$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

= \langle Base case, $n = 1$ \rangle

$$F_1 \leq \phi^{1-1}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

= $\langle \text{Base case, } n = 1 \rangle$

$$F_1 \leq \phi^{1-1}$$

= $\langle (12.14) \rangle$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

= $\langle \text{Base case, } n = 1 \rangle$

$$F_1 \leq \phi^{1-1}$$

= $\langle (12.14) \rangle$

$$1 \leq \phi^{1-1}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

= $\langle \text{Base case, } n = 1 \rangle$

$$F_1 \leq \phi^{1-1}$$

= $\langle (12.14) \rangle$

$$1 \leq \phi^{1-1}$$

= $\langle \text{Math} \rangle$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$F_n \leq \phi^{n-1}$$

= $\langle \text{Base case, } n = 1 \rangle$

$$F_1 \leq \phi^{1-1}$$

= $\langle (12.14) \rangle$

$$1 \leq \phi^{1-1}$$

= $\langle \text{Math} \rangle$

$$1 \leq 1 \quad //$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$F_n \leq \phi^{n-1}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \\ & F_2 \leq \phi^{2-1} \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \\ & F_2 \leq \phi^{2-1} \\ = & \langle (12.14) \text{ and math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \\ & F_2 \leq \phi^{2-1} \\ = & \langle (12.14) \text{ and math} \rangle \\ & 1 + 0 \leq \phi \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \\ & F_2 \leq \phi^{2-1} \\ = & \langle (12.14) \text{ and math} \rangle \\ & 1 + 0 \leq \phi \\ = & \langle (12.14.1) \text{ and math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Proof

First base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 1 \rangle \\ & F_1 \leq \phi^{1-1} \\ = & \langle (12.14) \rangle \\ & 1 \leq \phi^{1-1} \\ = & \langle \text{Math} \rangle \\ & 1 \leq 1 \quad // \end{aligned}$$

Second base case

$$\begin{aligned} & F_n \leq \phi^{n-1} \\ = & \langle \text{Base case, } n = 2 \rangle \\ & F_2 \leq \phi^{2-1} \\ = & \langle (12.14) \text{ and math} \rangle \\ & 1 + 0 \leq \phi \\ = & \langle (12.14.1) \text{ and math} \rangle \\ & 1 \leq 1.618 \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$F_{n+1}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \\ & \phi^{n-2}(\phi + 1) \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \\ & \phi^{n-2}(\phi + 1) \\ = & \langle (12.15) \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \\ & \phi^{n-2}(\phi + 1) \\ = & \langle (12.15) \rangle \\ & \phi^{n-2} \cdot \phi^2 \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \\ & \phi^{n-2}(\phi + 1) \\ = & \langle (12.15) \rangle \\ & \phi^{n-2} \cdot \phi^2 \\ = & \langle \text{Math} \rangle \end{aligned}$$

A Logical Approach to Discrete Math

Prove (12.16) $F_n \leq \phi^{n-1}$ for $n \geq 1$

Induction case

Prove $F_{n+1} \leq \phi^{(n+1)-1}$

assuming $F_n \leq \phi^{n-1}$ and $F_{n-1} \leq \phi^{(n-1)-1}$

as the inductive hypotheses.

$$\begin{aligned} & F_{n+1} \\ = & \langle (12.14) \text{ with } n := n + 1 \rangle \\ & F_n + F_{n-1} \\ \leq & \langle \text{Inductive hypotheses} \rangle \\ & \phi^{n-1} + \phi^{n-2} \\ = & \langle \text{Math, factor out } \phi^{n-2} \rangle \\ & \phi^{n-2}(\phi + 1) \\ = & \langle (12.15) \rangle \\ & \phi^{n-2} \cdot \phi^2 \\ = & \langle \text{Math} \rangle \\ & \phi^{(n+1)-1} \quad // \end{aligned}$$

A Logical Approach to Discrete Math

Inductively defined binary trees.

(12.30) **Definition, Binary Tree:**

\emptyset is a binary tree, called the empty tree.

(d, l, r) is a binary tree, for $d: \mathbb{Z}$ and l, r binary trees.



A Logical Approach to Discrete Math

Inductively defined binary trees.

(12.30) **Definition, Binary Tree:**

\emptyset is a binary tree, called the empty tree.

(d, l, r) is a binary tree, for $d: \mathbb{Z}$ and l, r binary trees.

⑤

$(5, \emptyset, \emptyset)$


A Logical Approach to Discrete Math

Inductively defined binary trees.

(12.30) **Definition, Binary Tree:**

\emptyset is a binary tree, called the empty tree.

(d, l, r) is a binary tree, for $d: \mathbb{Z}$ and l, r binary trees.



$(5, \emptyset, (3, \emptyset, \emptyset))$


A Logical Approach to Discrete Math

Inductively defined binary trees.

(12.30) **Definition, Binary Tree:**

\emptyset is a binary tree, called the empty tree.

(d, l, r) is a binary tree, for $d \in \mathbb{Z}$ and l, r binary trees.



$(5, (12, \emptyset, \emptyset), (3, (4, \emptyset, \emptyset), \emptyset))$

A Logical Approach to Discrete Math

(12.31) **Definition, Number of Nodes:**

$$\#\emptyset = 0$$

$$\#(d, l, r) = 1 + \#l + \#r$$

(12.32) **Definition, Height:**

$$\text{height}.\emptyset = 0$$

$$\text{height}.(d, l, r) = 1 + \max(\text{height}.l, \text{height}.r)$$

(12.32.1) **Definition, Leaf:** A leaf is a node with no children (i.e. two empty subtrees).

(12.32.2) **Definition, Internal node:** An internal node is a node that is not a leaf.

(12.32.3) **Definition, Complete:** A binary tree is complete if every node has either 0 or 2 children.

⑤

(5, \emptyset , \emptyset)

A Logical Approach to Discrete Math

(12.31) **Definition, Number of Nodes:**

$$\#\emptyset = 0$$

$$\#(d, l, r) = 1 + \#l + \#r$$

(12.32) **Definition, Height:**


$$\text{height}.\emptyset = 0$$

$$\text{height}.(d, l, r) = 1 + \max(\text{height}.l, \text{height}.r)$$

(12.32.1) **Definition, Leaf:** A leaf is a node with no children (i.e. two empty subtrees).

(12.32.2) **Definition, Internal node:** An internal node is a node that is not a leaf.

(12.32.3) **Definition, Complete:** A binary tree is complete if every node has either 0 or 2 children.



$(5, \emptyset, (3, \emptyset, \emptyset))$

A Logical Approach to Discrete Math

(12.31) **Definition, Number of Nodes:**

$$\#\emptyset = 0$$

$$\#(d, l, r) = 1 + \#l + \#r$$

(12.32) **Definition, Height:**


$$\text{height}.\emptyset = 0$$

$$\text{height}.(d, l, r) = 1 + \max(\text{height}.l, \text{height}.r)$$

(12.32.1) **Definition, Leaf:** A leaf is a node with no children (i.e. two empty subtrees).

(12.32.2) **Definition, Internal node:** An internal node is a node that is not a leaf.

(12.32.3) **Definition, Complete:** A binary tree is complete if every node has either 0 or 2 children.



$(5, (12, \emptyset, \emptyset), (3, (4, \emptyset, \emptyset), \emptyset))$

A Logical Approach to Discrete Math

- (12.33) The maximum number of nodes in a tree with height n is $2^n - 1$ for $n \geq 0$.
- (12.34) The minimum number of nodes in a tree with height n is $n + 1$ for $n \geq 0$.
- (12.35) (a) The maximum number of leaves in a tree with height n is 2^{n-1} for $n > 0$.
(b) The maximum number of internal nodes is $2^{n-1} - 1$ for $n > 0$.
- (12.36) (a) The minimum number of leaves in a tree with height n is 1 for $n > 0$.
(b) The minimum number of internal nodes is $n - 1$ for $n > 0$.
- (12.37) Every nonempty complete tree has an odd number of nodes.

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Base case

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Base case

(a) The empty tree has zero nodes.

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Base case

(a) The empty tree has zero nodes.

(b) $2^0 - 1 = 0$ //

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Induction case

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Induction case

Must prove “the maximum number of nodes in a tree of height $n+1$ is $2^{n+1} - 1$ ”

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Induction case

Must prove “the maximum number of nodes in a tree of height $n+1$ is $2^{n+1} - 1$ ”

assuming

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Induction case

Must prove “the maximum number of nodes in a tree of height $n+1$ is $2^{n+1} - 1$ ”

assuming

“the maximum number of nodes in a tree of height n is $2^n - 1$ ” as the inductive hypothesis.

A Logical Approach to Discrete Math

Prove (12.33) the maximum number of nodes in a tree of height n is $2^n - 1$.

Induction case


Must prove “the maximum number of nodes in a tree of height $n+1$ is $2^{n+1} - 1$ ”

assuming

“the maximum number of nodes in a tree of height n is $2^n - 1$ ” as the inductive hypothesis.


Proof: A tree height $n+1$ with the maximum number of nodes must have two children of height n , each with the maximum number of nodes.

A Logical Approach to Discrete Math




A Logical Approach to Discrete Math

By the inductive hypothesis




A Logical Approach to Discrete Math

By the inductive hypothesis
I has $2^n - 1$ nodes




A Logical Approach to Discrete Math

By the inductive hypothesis
l has $2^n - 1$ nodes
and r has $2^n - 1$ nodes.




A Logical Approach to Discrete Math

By the inductive hypothesis
l has $2^n - 1$ nodes
and r has $2^n - 1$ nodes.
So, including the root node,



A Logical Approach to Discrete Math


By the inductive hypothesis
l has $2^n - 1$ nodes
and r has $2^n - 1$ nodes.
So, including the root node,
the total is



A Logical Approach to Discrete Math

By the inductive hypothesis
l has $2^n - 1$ nodes
and r has $2^n - 1$ nodes.
So, including the root node,
the total is

$$1 + (\# \text{ in } l) + (\# \text{ in } r)$$




A Logical Approach to Discrete Math

By the inductive hypothesis
l has $2^n - 1$ nodes
and r has $2^n - 1$ nodes.

So, including the root node,
the total is

$$= 1 + (\# \text{ in } l) + (\# \text{ in } r)$$


<Ind. hyp.>



A Logical Approach to Discrete Math

By the inductive hypothesis
l has 2^{n-1} nodes
and r has 2^{n-1} nodes.
So, including the root node,
the total is


$$\begin{aligned} & 1 + (\# \text{ in } l) + (\# \text{ in } r) \\ = & \quad \langle \text{Ind. hyp.} \rangle \\ & 1 + 2^{n-1} + 2^{n-1} \end{aligned}$$



A Logical Approach to Discrete Math

By the inductive hypothesis
l has 2^{n-1} nodes
and r has 2^{n-1} nodes.
So, including the root node,
the total is


$$\begin{aligned} & 1 + (\# \text{ in } l) + (\# \text{ in } r) \\ = & \quad \langle \text{Ind. hyp.} \rangle \\ & 1 + 2^{n-1} + 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \end{aligned}$$



A Logical Approach to Discrete Math

By the inductive hypothesis
l has 2^{n-1} nodes
and r has 2^{n-1} nodes.
So, including the root node,
the total is


$$\begin{aligned} & 1 + (\# \text{ in } l) + (\# \text{ in } r) \\ = & \quad \langle \text{Ind. hyp.} \rangle \\ & 1 + 2^{n-1} + 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \\ & 1 + 2 \times 2^{n-1} \\ & 1 + 2^n - 2 \end{aligned}$$



A Logical Approach to Discrete Math

By the inductive hypothesis
l has 2^{n-1} nodes
and r has 2^{n-1} nodes.
So, including the root node,
the total is

$$\begin{aligned} & 1 + (\# \text{ in } l) + (\# \text{ in } r) \\ = & \quad \langle \text{Ind. hyp.} \rangle \\ & 1 + 2^{n-1} + 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \\ & 1 + 2 \times 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \end{aligned}$$



A Logical Approach to Discrete Math

By the inductive hypothesis
l has 2^{n-1} nodes
and r has 2^{n-1} nodes.
So, including the root node,
the total is

$$\begin{aligned} & 1 + (\# \text{ in } l) + (\# \text{ in } r) \\ = & \quad \langle \text{Ind. hyp.} \rangle \\ & 1 + 2^{n-1} + 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \\ & 1 + 2 \times 2^{n-1} \\ = & \quad \langle \text{Math} \rangle \\ & 2^{n+1} - 1 \quad // \end{aligned}$$

